## **NEW PATIENT NOTES – RECALL SET UP**

FEMALES	MALES
CVD	CVD
45 year vaccine	45 year vaccine
65 year vaccines	65 year vaccines
-Boostrix	-Boostrix
-Shingles	-Shingles
-Flu	-Flu
Childhood vaccines	Childhood vaccines
11 year vaccine	11 year vaccine
HPV vaccines	HPV vaccines
If diabetic – recalls	If diabetic – recalls
Smears	
Mammograms	

- Doing NIR search on children to check status
- Doing NIR to check if had HPV vaccines up to 27 years
- Getting smear history from the cervical screening register
- If new to the Waikato checking if they want to be enrolled with Waikato Breast Screening.

## Table 4: When to start risk assessments for men and women in different population subgroups

Population subgroup	Men	Women
Individuals without known risk factors	Age 45 years	Age 55 years
Māori, Pacific peoples or South-Asian* peoples	Age 30 years	Age 40 years
People with other known cardiovascular risk factors or at high risk of developing diabetes	Age 35 years	Age 45 years

## Family history risk factors:

- · diabetes in first-degree relative (parent, brother or sister)
- hospitalisation for or death from heart attack or stroke in a firstdegree relative before the age of 50 years (father or brother, mother or sister)
- Familial hypercholesterolaemia

## Personal history risk factors:

- · people who smoke
- gestational diabetes
- HbA<sub>1c</sub> 41–49 mmol/mol
- BMI ≥ 30 or truncal obesity (waist circumference ≥ 102 cm in men or ≥ 88 cm in women)
- eGFR<60 but >45 ml/min/1.73 m<sup>2</sup> †
- atrial fibrillation

People with diabetes (type 1 or 2)	From the time of diagnosis	From the time of diagnosis
People with severe mental illness (Appendix C)	From age 25 years	From age 25 years

South-Asian peoples: Indian, including Fijian Indian, Sri Lankan, Afghani, Bangladeshi, Nepalese, Pakistani, Tihetan

<sup>†</sup> eGFR estimated glomerular filtration rate.